

(7 February 2012)

DECONVOLUTION OF POINT PROCESSESDANIEL EDWARD CLARK,^{*}**Abstract**

The superposition of two independent point processes can be described by multiplication of their probability generating functionals (p.g.fl.s). The inverse operation, which can be viewed as a deconvolution, is defined by dividing the superposed process by one of its constituent p.g.fl.s. The deconvolved process is computed using the higher-order chain rule for Gâteaux differentials. The higher-order quotient rule for Gâteaux differentials is first established and then applied to point processes.

Keywords: point processes; generating functionals; Gâteaux differentials

2010 Mathematics Subject Classification: Primary 60G55

Secondary 49J50

The superposition of two point processes can be formed by multiplication of their probability generating functionals [6]. While it is clear that the inverse operation would involve division of the probability generating functionals, there are no results regarding the Janossy densities after this operation. This is probably due to the fact that the denominator involves an infinite sum and there was no easy way of computing the point process. In this article, I derive the deconvolved point process using the recently derived higher-order chain rule for Gâteaux differentials [1].

The paper is structured as follows. Section 1 describes the necessary background on functionals to derive the result, including the probability generating functional (p.g.fl.), the Gâteaux differential, Janossy densities and the higher-order chain rule. Section 2 presents the main result by first establishing the n^{th} -order Gâteaux differential of the reciprocal of a functional in Lemma 4. Theorem 1 establishes the higher-order quotient rule for Gâteaux differentials using Leibniz' rule and the recently derived Faa di Bruno's formula for Gâteaux differentials [1]. This is used to determine the expression for the deconvolved point process via the probability generating functional [6].

1. Probability generating functionals

This section discusses generating functionals as a means of describing systems with a variable number of objects. Gâteaux differentials are discussed as a means of determining their constituent functions. The higher-order chain rule is introduced to determine the differentials of composite functionals [1].

^{*} Postal address: School of Engineering and Physical Sciences, Heriot-Watt University, Edinburgh, EH14 4AS UK

Definition: probability generating functional

Following Moyal [6], define the *probability generating functional* (p.g.fl.) with

$$G_X(\psi) = p_0 + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\int \prod_{i=1}^n dx_i \psi(x_i) \right) p_n(x_1, \dots, x_n), \quad (1)$$

where $p_n(x_1, \dots, x_n) \geq 0$ and

$$p_0 + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\int \prod_{i=1}^n dx_i \right) p_n(x_1, \dots, x_n) = 1 \quad (2)$$

The p.g.fl. is commonly used in point process theory to characterise point patterns [2]. For compactness of notation, we write the p.g.fl. as

$$G_X(\psi) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\int \prod_{i=1}^n dx_i \psi(x_i) \right) p_n(x_1, \dots, x_n). \quad (3)$$

Definition: Gâteaux differential

A *Gâteaux differential* [4, p71-74], of a functional $\Psi(\psi)$ with increment ξ , can be defined with

$$\delta\Psi(\psi; \xi) = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} (\Psi(\psi + \epsilon\xi) - \Psi(\psi)). \quad (4)$$

The n^{th} -order differential (or variation), which is defined by recursively, is denoted $\delta^n\Psi(\psi, \xi_1, \dots, \xi_n)$.

Lemma 1

The p.g.fl. $G(h)$ defines P uniquely, since the Janossy densities can be recovered with Gâteaux differentials [3], i.e.

$$p_k(x_1, \dots, x_k) = \delta^k G(0; \xi_1, \dots, \xi_k), \quad (5)$$

where $\xi_i(x) = \delta_{x_i}(x)$.

Proof

See Moyal [6].

Lemma 2

Let Π be the set of all partitions of variables $\eta_1, \dots, \eta_n$, and $\pi \in \Pi$ denote a single partition that has constituent blocks $\omega \in \pi$ of size $|\omega|$ consisting of constituent elements $\zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|} \in \{\eta_1, \dots, \eta_n\}$. The n^{th} -order variation of composition $f \circ g$ with increments $\eta_1, \dots, \eta_n$ is given by

$$\delta^n((f \circ g)(y); \eta_1, \dots, \eta_n) = \sum_{\pi \in \Pi} \delta^{|\pi|} f(g(y); \xi_{\pi, 1}(y), \dots, \xi_{\pi, |\pi|}(y)), \quad (6)$$

where $\xi_{\pi, \omega}(y)$ is the variation of order $|\omega|$ with increments $\zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|}$, i.e.

$$\xi_{\pi, \omega}(y) = \delta^{|\omega|} g(y; \zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|}). \quad (7)$$

Proof

See Clark [1].

Lemma 3

Leibniz' rule for functional derivatives the differentials of the product $f(y)g(y)$ with increments $\Xi = \{\eta_1, \dots, \eta_n\}$ is

$$\delta^n (f(y)g(y); \eta_1, \dots, \eta_n) = \sum_{\Phi \subset \Xi} \delta^{|\Phi|} f(y; \phi_1, \dots, \phi_{|\Phi|}) \delta^{|\Xi| - |\Phi|} g(y; \xi_1, \dots, \xi_{|\Xi| - |\Phi|}), \quad (8)$$

where the summation is over all subsets Φ of the increments, and $\Phi = \{\phi_1, \dots, \phi_{|\Phi|}\}$, $\Xi - \Phi = \{\xi_1, \dots, \xi_{|\Xi| - |\Phi|}\}$.

Proof

See Mahler [5, p389].

2. Deconvolution of point processes

This section presents the main result. Lemma 4 establishes the higher-order differential of the reciprocal of a functional, which is then used to establish a formula for the higher-order quotient rule for Gâteaux differentials in Theorem 1. It is then shown in that the result is directly applicable to point processes.

Lemma 4

Consider a functional $g(y)$ in Banach space χ . Then the n^{th} -order Gâteaux differential [3] of $1/g(y)$ with set of increments $\Pi = \{\eta_1, \dots, \eta_n\}$ is found with

$$\delta^n (1/g(y); \eta_1, \dots, \eta_n) = \sum_{\pi \in \Pi} \frac{(-1)^{|\pi|} |\pi|!}{g(y)^{|\pi|+1}} \prod_{\omega \in \pi} \delta^{|\omega|} g(y; \zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|}), \quad (9)$$

where the summation is over all partitions π of increment set Π . The product is over all cells $\omega = \{\zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|}\}$ of partition π .

Proof

Suppose that $f(x) = 1/x$, where x is real-valued and non-zero. Then, following the definition of the Gâteaux derivative (which in this case reduces to the same rules as ordinary derivative),

$$\delta^n (1/x; \eta_1, \dots, \eta_n) = \frac{(-1)^n n!}{x^{n+1}}. \quad (10)$$

Substituting this into (6) gives the required result.

Theorem 1

The higher-order quotient rule for Gâteaux differentials with increments $\Xi = \{\eta_1, \dots, \eta_n\}$ is determined with

$$\delta^n \left(\frac{f(y)}{g(y)}; \eta_1, \dots, \eta_n \right) = \sum_{\Pi \subset \Xi} \left(\sum_{\pi \in \Pi} \frac{(-1)^{|\pi|} |\pi|!}{g(y)^{|\pi|+1}} \prod_{\omega \in \pi} \delta^{|\omega|} g(y; \zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|}) \right) \delta^{|\Xi| - |\Pi|} f(y; \phi_1, \dots, \phi_{|\Xi| - |\Pi|}), \quad (11)$$

where the summation on the left is over all subsets Π of Ξ . The inner summation is over all partitions π of subset Π . The term on the right, $\delta^{|\omega|} g(y; \zeta_{\pi, \omega, 1}, \dots, \zeta_{\pi, \omega, |\omega|})$ is the variation of order $|\Xi| - |\Pi|$ with increments $\phi_1, \dots, \phi_{|\Xi| - |\Pi|}$ that comprise $\Xi - \Pi$.

Proof

The result follows from application of Lemmas 3 and 4.

Corollary

Consider a point process defined by the superposition of point processes with p.g.f.s, i.e.

$$G[h] = G_1[h]G_2[h], \quad (12)$$

defined with Janossy densities $p(x_1, \dots, x_n)$, for $G[h]$, $q(x_1, \dots, x_n)$, for $G_1[h]$, and $r(x_1, \dots, x_n)$, for $G_2[h]$, for $n > 0$. Let $\Xi = \{x_1, \dots, x_n\}$. Then we can recover the Janossy densities $r(x_1, \dots, x_n)$ of $G_2[h]$ with

$$r(x_1, \dots, x_n) = \sum_{\Pi \subset \Xi} \left(\sum_{\pi \in \Pi} \frac{(-1)^{|\pi|} |\pi|!}{q_0^{|\pi|+1}} \prod_{\omega \in \pi} q(v_{\pi, \omega, 1}, \dots, v_{\pi, \omega, |\omega|}) \right) p(w_1, \dots, w_{|\Xi| - |\Pi|}), \quad (13)$$

where the summation on the left is over all subsets Π of Ξ . The inner summation is over all partitions π of subset Π . The term on the right $p(w_1, \dots, w_{|\Xi| - |\Pi|})$ is the Janossy density evaluated at $w_1, \dots, w_{|\Xi| - |\Pi|}$ which comprise subset $\Xi - \Pi$, and q_0 is the zeroth term in $G_1[h]$. When $n = 0$, $r_0 = p_0/q_0$.

Proof

Let $G_1[h] = f(y)$ and $G_2[h] = g(y)$ in (i). Using Lemma 1, and setting $\eta_1 = \delta_{x_1}, \dots, \eta_n = \delta_{x_n}$, and $h = 0$ yields the result for $n > 0$. For $n = 0$, we note that $G[0] = p_0$, and $G_1[0] = q_0$, hence $r_0 = p_0/q_0$.

References

- [1] CLARK, D. E. (2012). Faà di Bruno's formula for Gâteaux differentials. *arXiv:1202.0264*.
- [2] COX, D. R. AND ISHAM, V. (1980). *Point Processes*. Chapman and Hall.
- [3] GÂTEAUX, R. (1919). Fonctions d'une infinité de variables indépendantes. *Bulletin de la Société Mathématique de France* **47**, 70–96.
- [4] HILLE, E. (1948). Functional Analysis and Semi-Groups. *Amer. Math. Soc.*

- [5] MAHLER, R. (2007). Statistical Multisource Multitarget Information Fusion. *Artech House*.
- [6] MOYAL, J. E. (1962). The General Theory of Stochastic Population Processes. *Acta Mathematica* **108**, 1–31.